

<https://helda.helsinki.fi>

Presence of Chlamydophila pneumoniae DNA in blood cells is a frequent event in patients with the late stage of primary cutaneous lymphomas and with atopic dermatitis

Nedoszytko, Boguslaw

2018

Nedoszytko , B , Wierzbicki , P , Karenko , L , Maciejewska-Radomska , A , Stachewicz , P , Zablotna , M , Glen , J , Vakeva , L , Nowicki , R J & Sokolowska-Wojdylo , M 2018 , ' Presence of Chlamydophila pneumoniae DNA in blood cells is a frequent event in patients with the late stage of primary cutaneous lymphomas and with atopic dermatitis ' , Postepy Dermatologii i Alergologii , vol. 35 , no. 3 , pp. 274-279 . <https://doi.org/10.5114/ada.2018.76224>

<http://hdl.handle.net/10138/237143>

<https://doi.org/10.5114/ada.2018.76224>

unspecified

publishedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

Presence of *Chlamydomphila pneumoniae* DNA in blood cells is a frequent event in patients with the late stage of primary cutaneous lymphomas and with atopic dermatitis

Bogusław Nedoszytko¹, Piotr Wierzbicki², Leena Karenko³, Agata Maciejewska-Radomska⁴, Przemysław Stachewicz², Monika Zabłotna¹, Jolanta Gleń¹, Liisa Väkevää³, Roman J. Nowicki¹, Małgorzata Sokołowska-Wojdyło¹

¹Department of Dermatology, Venereology and Allergology, Medical University of Gdansk, Gdansk, Poland

²Department of Histology, Medical University of Gdansk, Gdansk, Poland

³Department of Dermatology and Allergology, University of Helsinki and Helsinki University Central Hospital, Helsinki, Finland

⁴Individual Specialist Medical Practice, Gdynia, Poland

Adv Dermatol Allergol 2018; XXXV (3): 274–279

DOI: <https://doi.org/10.5114/ada.2018.76224>

Abstract

Introduction: Microbial infection and associated super antigens have been implicated in the pathogenesis of cutaneous T-cell lymphoma (CTCL), and many patients die from complicating bacterial infections. It has been postulated that *Chlamydomphila pneumoniae* (*C. pneumoniae*) infection may be involved in the pathogenesis of *Mycosis fungoides* (MF) but published data are limited and controversial.

Aim: To analyze the frequency of (*C. pneumoniae*) DNA presence in blood samples of lymphoma cases.

Material and methods: Using Q-PCR method we analyzed the presence of DNA in the blood samples obtained from 57 patients with CTCL (55 – mycosis fungoides (MF)/Sézary syndrome (SS), one primary cutaneous anaplastic large cell lymphoma (CD30+) and one NKT cell lymphoma) and 3 patients with cutaneous B-cell lymphomas, and 120 individuals from control groups (40 patients with psoriasis, 40 patients with atopic dermatitis and 40 healthy controls).

Results: *Chlamydomphila pneumoniae* DNA was identified in 13 of 55 cases in the MF/SS group (23.6%), in 1 patient with CD30+ large cell lymphoma and in 1 of 3 patients with B-cell lymphoma. The presence of *C. pneumoniae* was confirmed in 1 of 40 psoriatic patients (2.5%), in 5 of 40 patients with atopic dermatitis (12.5%) and in none of 40 healthy individuals. Presence of *C. pneumoniae* DNA in MF patients was strongly associated with disease progression; $r_s = 0.756$; $p = 0.0123$ for groups IA → IVB, and was noted more frequently in advanced (III + IV) stages than in early (I–II) stages ($p = 0.0139$). There are no differences in the mean age of MF/SS patients with and without infection.

Conclusions: The presence of *C. pneumoniae* DNA in the blood cells is a frequent event in late stages of MF/SS and may be explained by Th2 shift and suppression of the immune system during the course of the disease.

Key words: *Chlamydomphila pneumoniae*, late mycosis fungoides/Sézary syndrome stages, polymerase chain reaction.

Introduction

Antigen stimulation by pathogens such as bacteria and viruses has been considered a possible predisposing factor to uncontrolled cell proliferation and development of lymphoid and other neoplastic tissue. *Helicobacter pylori* precedes the development of gastric B-cell lymphoma, mycoplasma-like organisms and hepatitis C virus have been suggested to be associated with Hodgkin disease

and *Chlamydia trachomatis* infection may associate with rectal and cervical cancer. Although several etiologies have been postulated for mycosis fungoides (MF) and Sézary syndrome (SS), their causes remain unknown. Based on the limited experimental and clinical data, it has been speculated that chronic local antigen stimulation by *Staphylococcus aureus* as well as *Chlamydomphila pneumoniae* (*C. pneumoniae*) may play a role in the pathogenesis of cutaneous T-cell lymphoma (CTCL) [1–3]. *Chlamydo-*

Address for correspondence: Małgorzata Sokołowska-Wojdyło MD, PhD, Department of Dermatology, Venereology and Allergology, Medical University of Gdansk, 7 Dębinki St, 80-210 Gdansk, Poland, phone: +48 602 618 195, e-mail: mwojd@gumed.edu.pl

Received: 1.02.2017, **accepted:** 14.04.2017.

phila pneumoniae is a common intracellular micro-organism. Seroepidemiological studies indicate that *C. pneumoniae* infection is by far the most common human chlamydial infection in different cohorts, with seropositivity in at least 50% of the general population over the age of 20 [4–10]. But polymerase chain reaction (PCR) studies on asymptomatic healthy adults (more than 1000) had established only 1% of positivity in nasopharyngeal swabs' specimens [11]. In addition to pneumonia, pharyngitis, bronchitis and asthma *C. pneumoniae* is also associated with arteriosclerosis, lung cancer, multiple sclerosis and Alzheimer's disease [12–18]. *Chlamydophila pneumoniae* can infect, reside and replicate in various cell types including smooth muscle cells, fibroblasts, endothelial cells, bronchial epithelial cells, keratinocytes as well as various immune cells such as macrophages, lymphocytes and natural killer cells (NK) [19, 20]. It induces the increased release of pro-inflammatory mediators including tumor necrosis factor α (TNF- α), interleukin 6 and 8 (IL-6, IL-8), basic fibroblast growth factor (bFGF) and up regulates adhesion molecules [21]. Recently it has been suggested that *C. pneumoniae* infection may also stimulate the IL-10 production which down regulates the expression of major histocompatibility complex class I (MHC-I), inhibits apoptosis and increases the longevity of the host cell, enhancing the survival of bacteria itself [22–24]. The role of *C. pneumoniae* in the etiology of CTCL is controversial. It has been suspected that a localized bacterial infection increases local production of inflammatory cytokines including interferon γ (IFN- γ) (critical in immunity and immunopathology of chlamydial infection) and CXCL-10, a cytokine chemo attractive for epidermotropic T lymphocytes [24]. Studies on the growth requirements of the abnormal T lymphocytes in MF/SS lead to the identification of a so-called Sézary cell activation factor (SAF) that stimulates the growth of both malignant and non-malignant T cells. Sézary cell activation factor was originally defined as an inducer of functional IL-2 receptors. It is postulated that a combination of SAF and IL-2 stimulates the propagation of oligoclonal T-cell populations from the peripheral blood mononuclear cells (PBMC) of patients with SS, with approximately one third of those cell clones containing the predominant malignant clone [3, 25]. Using a monoclonal antibody inhibitory for SAF activity Abrams *et al.* demonstrated that SAF is present in more than half skin biopsies taken from patients with MF [26]. It was also confirmed that SAF determinant is not of eukaryotic origin and is associated with *C. pneumoniae* bacteria [3]. Sézary cell activation factor is a protein of approximately 30 kDa, resembling the *C. pneumoniae* T cell activation factor originally described by Halme *et al.* [27]. Abrams *et al.* confirmed the presence of *C. pneumoniae* DNA and RNA in the skin by PCR and reverse transcription – PCR and by sequence analysis of the PCR products. The authors showed that *C. pneumoniae* antigen expression was associated

with active disease and was not found after psoralen and ultraviolet therapy.

Material and methods

Study and control groups

In this study 60 patients with skin lymphomas (48 from Poland – 30 men and 18 women, mean age: 60.7 \pm 13.5 and 12 from Finland – 8 men and 4 woman, mean age: 61.9 \pm 21.6) were included, of whom 57 patients were diagnosed with primary CTCL and 3 patients with primary cutaneous B-cell lymphomas (one follicle-center and two marginal zone lymphomas) according to WHO criteria. The control groups consisted of 40 patients with psoriasis (22 men and 18 women, mean age: 49.3 \pm 14.3), 40 with atopic dermatitis (21 men and 19 women, mean age: 13.8 \pm 7.7) and 40 healthy individuals (21 men, 19 women, 10 from Finland and 30 from Poland, mean age: 39 \pm 15.0). From the 57 CTCL patients, 55 have MF or SS, 1 patient had primary cutaneous anaplastic large cell lymphoma CD (30+), and one had NK/T cell lymphoma. From 55 MF/SS patients, 20 were in early clinical stages (IA–IIA) and 35 patients were in advanced stages (IIB–IVB). Additionally skin biopsies were obtained from 6 MF/SS patients.

Material collection and DNA samples

From all patients and control groups peripheral whole blood samples (EDTA-K2 (Medlab Products, Raszyn, Poland) were collected and stored at -80°C . Additionally from 6 examined patients skin biopsies were collected and stored at -80°C . DNA was extracted using Blood Mini or Mini AX Tissue (A&A Biotechnology, Gdynia, Poland) according to the manufacturer's protocols. Further, DNA was precipitated using common sodium acetate-ethanol technique [1] in a final volume of 20 μl and stored at -20°C for further analyses. DNA concentration and purity were assessed by NanoDrop ND1000 (ThermoScientific, Wilmington, DE, USA).

Chlamydophila pneumoniae strain and cell lines of *C. pneumoniae* TW183 strain was purchased from American Type Culture Collection (ATCC) (Rockville, MD, USA) and stored at -80°C . In order to propagate *C. pneumoniae* for PCR optimization, HELA 229 line (ATCC) cells were infected by TW183 lysate in a 24-cell plate, followed by centrifugation (15 min, 400 \times G) and incubated at 37°C and 5% of CO_2 for 2 days, according to ATCC protocol. After incubation, cells were harvested and stored at -80°C . Presence of *C. pneumoniae* in cells was further confirmed by diagnostic nested-PCR assay (BLIRT-DNA, Gdansk, Poland).

Quantitative polymerase chain reaction analysis

In order to create a specific quantitative assay for *C. pneumoniae* detection we checked (BLAST N database)

15 reference sequences of the *C. pneumoniae* strains and for diversification *C. trachomatis* strains and the ompA gene was chosen as a good molecular target for detection and quantification.

Optimization and validation of QPCR assay

We amplified 805 bp fragment of *C. pneumoniae* TW183 strain based on following primers: 5'CCGGCCTA-CAATAAGCATTTAC and 5'GAGCTTCTGCAGTAAGTGACCA. The sequence was confirmed by RFLP method. After purification, the PCR product was cloned into pUC19 plas-

mid, followed by propagation in *Escherichia coli* strain, isolation, purification and spectrophotometric quantification of plasmid. This construct was applied as a positive control for the calibration curve. The primers for QPCR assay were designed using VNTI software (Invitrogen, Life Technologies, Carlsbad, CA, USA): 5'AACAAAGTCTGCGAC-CATCAATTAC, 5'GGCTGAGCAATGCGGATGTTATCAC. The conditions of PCR reaction: 2× SybrGreen Supermix (Bio-Rad, Hercules, CA, USA), 170 nM of each primer, 4 mM MgCl₂ and ddH₂O were mixed with 2 µl of template DNA to a final volume of 17 µl. QPCR was performed in iCycler

Table 1. Frequency of *Chlamydomphila pneumoniae* DNA in blood samples from patients with various types of lymphomas, psoriasis, atopic dermatitis and healthy control groups

Diagnosis/clinical stage	Frequency of <i>Chlamydomphila pneumoniae</i> DNA			I vs. II vs. III–IV stages	Early vs. late stage
	Polish patients (<i>n</i> = 48)	Finnish patients (<i>n</i> = 12)	Both patient groups (<i>n</i> = 60)		
MF patients:					
IA	0/2	0/1	0/3 (0%)	0/16 (0%)	1/20 (5.0%) <i>p</i> = 0.0139
IB	0/13	0/0	0/13 (0%)		
IIA	1/3	0/1	1/4 (25%)	3/17 (17.6%)	
IIB	1/11	1/2	2/13 (15%)		
III	3/6	0/0	3/6 (50%)	6/11 (54.5%)	
IVA	0/0	1/3*	1/3 (33%)		
IVB	2/2	0/0	2/2 (100%)		
MF summary			9/44 (20.5%)		
SS patients:					
IVA	0/1	0/0	0/1		
IVB	4/5	0/5	4/10 (40%)		
SS summary			4/11 (36.4%) (MF vs. SS, NS, <i>p</i> = 0.266)		
All MF/SS patients	11/43 (25.6%)	2/12 (16.7%)	13/55 (23.6%)		
c-ALCL (CD30+)	1/1	0	1/1		
NK/T lymphoma	0/1	0	0/1		
All CTCL patients			14/57 (24.5%)		
Primary cutaneous B-cell lymphomas	1/3	0	1/3 (33%)		
All lymphomas			15/60 (25.0%)		
Controls					
AD patients					5/40 (12.5%)
Psoriasis patients					1/40 (2.5%)
Healthy individuals					0/40 (0%)

Positive results obtained only from skin biopsy material. MF – mycosis fungoides, SS – Sezary syndrome, c-ALCL – cutaneous anaplastic large cell lymphoma, AD – atopic dermatitis.

and fluorescence data were automatically collected and analyzed by iCycler iQ Optical Software ver. 3.0a (Bio-Rad, Hercules, CA, USA). QPCR conditions were: initial denaturation – 95°C/3 min; 40× (95°C/20 s, 65°C/30 s, 72°C/20 s, 77°C/5 s-fluorescence reading step). A calibration curve was performed in triplicates using 6 ten-fold dilutions (10^{-8} to 10^{-3}) of pUC19 plasmid DNA containing ompA gene. The slope was -3.497 ; efficiency = 93.2% with a correlation coefficient of $R^2 = 0.997$. Dynamic melt-curve analysis and agarose-gel electrophoresis were used for all post-PCR reaction tubes to confirm the size of expected amplicon (147 bp). All reactions containing analyzed DNA were performed in duplicates. If the ΔCt between replicates was > 0.3 and/or we found a melting peak with a different melting point $> 1^\circ\text{C}$ from the expected one, the reactions were repeated. As a method of quantification we chose the relative quantity method and for histograms we applied ΔCt method, where $\Delta Ct = (\text{mean } Ct \text{ of analyzed group}) - (\text{mean } Ct \text{ of calibration points})$.

Statistical analysis

All statistical analyses were done using the Statistica 8.0 (StatSoft Inc., Tulsa, OK, USA), Pearson and Fisher's exact test. For all statistical tests, we used a comparison related significance level of $p < 0.05$.

Results

The presence of *C. pneumoniae* DNA was detected in 14 of 57 (24.5%) CTCL patients and in 1 of 3 patients with B-cell lymphomas (primary cutaneous follicle-center B cell lymphoma). In the CTCL group, positive results were found in 13 of 55 (23.6%) MF/SS patients and in one patient with CD30 (+) anaplastic large cell lymphoma. *Chlamydophila pneumoniae* was detected only in 1 of 6 analyzed skin biopsies taken from MF/SS patients (Table 1). From the controls, bacterial DNA was present in the blood samples in 1 of 40 (2.5%) psoriasis patients, in 5 of 40 (12.5%) patients with atopic dermatitis, and in none of 40 healthy individuals. No differences between MF and SS patients were observed ($p = 0.266$). The frequency *C. pneumoniae* DNA in the MF/SS group correlated with disease progression ($rs = 0.756$, $p = 0.0123$, Figure 1). None of 16 samples from MF patients with disease (IA or IB) were positive while *C. pneumoniae* was detected in 3 of 17 (17.6%) cases in stage IIA or IIB, and in 6 of 11 (54.5%) MF patients in stages III or IV of disease. In SS patients, *C. pneumoniae* DNA was found in 4 of 11 (36.4%) patients with stage IVB. Cumulatively in the MF/SS group, *C. pneumoniae* infection was found in 1 of 20 (5.0%) patients in early clinical stage (IA–IIA) and in 12 of 35 (34.3%) patients in advanced clinical stage (IIB–IVB). The results were statistically significant ($p = 0.0139$). No differences in mean age between MF/SS patients with and without infection were observed (Figure 1).




Figure 1. Frequency of *Chlamydophila pneumoniae* in mycosis fungoides/Sézary syndrome patients with different clinical stage of disease

Discussion

Cutaneous T-cell lymphoma is a malignancy of skin-homing Th1 and/or Th₂ T cells. A prominent feature of CTCL is immunosuppression, which increases the risk of bacterial and viral infections in patients especially in the advanced stages of disease. The pathophysiology of this immunodeficiency is probably multifactorial. Data from experimental studies suggest that the T-cell repertoire in CTCL patients is significantly contracted [27]. Immunological abnormalities in CTCL are typically associated with depressed ability of peripheral blood cells to produce the Th1 cytokines, IFN- γ and IL-2 as a result of Th2 skewing [28–37]. During CTCL progression, reduced T-cell-mediated cellular immune responses and diminished natural killer cell activity also develop [38, 39].

Wysocka *et al.* have demonstrated a direct relationship between the extent of the pool of circulating malignant T-cells and an impaired immune response [40]. Also the functions of natural killer (NK) cells, including cellular cytotoxicity and production of IFN- γ , become increasingly impaired as the circulating tumor burden increases. The authors showed an inverse correlation between circulating clonal T cells and activation status of both NK cells and CD8 T cells with a diminishing expression of a number of the activation markers CD69 and CD25, as well as decreased intracellular IFN- γ production [40]. The following observations have two main pathophysiological implications. The impaired cellular immune response, which is pivotal for direct antitumor responses, leads to further acceleration of growth of the malignant T-cells population. Another consequence of the decline in cytotoxic T-cell and NK cell functions is impaired activity against opportunistic infectious pathogens. This theory has been supported by the clinical observations [41–43]. It is suggested that the Th2 cytokine pattern may create a permissive environment for *C. pneumoniae* infection. This concept is supported by few experimental studies

and clinical observations [44]. Referring to our results, *C. pneumoniae* DNA was found in 25% of all patients with CTCL. The frequency of chlamydial infection correlated with the stage of disease. The positive results were seen in 5% and 34% of patients with low and high clinical stages of disease, respectively. This observation may confirm the important role of impaired cellular immunological response in the control of opportunistic infections such as *C. pneumoniae* in CTCL patients. The association between Th1 response impairment and *C. pneumoniae* infection may be also supported by an increased incidence of bacterial DNA in patients with atopic dermatitis (12.5%) (Th2 cytokines pattern) vs. psoriatic patients (2.5%) (Th1 cytokines pattern) and healthy controls (0%) (immunocompetent individuals). *Chlamydomphila pneumoniae* DNA was detected only in one skin biopsy in our study group, in accordance with the results of German and Italian investigators, who did not detect any bacterial DNA in skin biopsies from MF/SS patients, and confirm that *C. pneumoniae* infection could not be estimated as a primary triggering factor in the pathogenesis of CTCL, in spite of Casselli *et al.* who had described skin presence of *Chlamydia* spp. (and HHV8) in all recurrences of CD30+CTCL as well as in routine control blood samples [45–47].

Conclusions

Our results suggest that *C. pneumoniae* infection is not a primary event in the pathogenesis of CTCL, but may be estimated as a risk factor complicating advanced stages of the disease associated with Th2 shift and deficiency of antibacterial defence mechanisms.

Acknowledgments

The study was approved by the local research ethics committee at the Medical University of Gdansk. The study is financed by the Polish Ministry of Science and Higher Education, grant 02-0066/07/253.

Conflict of interest

The authors declare no conflict of interest.

References

1. Tan RSH, Butterworth CM, McLaughlin H, et al. Mycosis fungoides: a disease of antigen persistence. *Br J Dermatol* 1974; 91: 607-16.
2. Jackow CM, Cather JC, Hearne V, et al. Association of erythrodermic cutaneous T-cell lymphoma, superantigen – positive *Staphylococcus aureus* and oligoclonal T-cell receptor V beta gene expansion. *Blood* 1997; 89: 32-40.
3. Abrams JT, Balin BJ, Vonderheid EC. Association between Sezary T-cell activating factor, *Chlamydia pneumoniae*, and cutaneous T cell lymphoma. *Ann N Y Acad Sci* 2001; 941: 69-85.
4. Kanamoto Y, Ouchi K, Mizui M, et al. Prevalence of antibody to *Chlamydia pneumoniae* TWAR in Japan. *J Clin Microbiol* 1991; 29: 816-8.
5. Marton A, Karolyi A, Szalka A. Prevalence of *Chlamydia pneumoniae* antibodies in Hungary. *Eur J Clin Microbiol Infect Dis* 1992; 11: 139-42.
6. Montes M, Cilla G, Alcorta M, et al. High prevalence of *Chlamydia pneumoniae* infection in children and young adults in Spain. *Pediatr Infect Dis J* 1992; 11: 972-3.
7. Koivisto AL, Isoaho R, Von Hertzen L, et al. Chlamydial antibodies in an elderly Finnish population. *Scand J Infect Dis* 1999; 31: 153-9.
8. Ni AP, Lin GY, Yang L, et al. A seroepidemiologic study of *Chlamydia pneumoniae*, *Chlamydia trachomatis* and *Chlamydia psittaci* in different populations on the mainland of China. *Scand J Infect Dis* 1996; 28: 553-7.
9. Ferrari M, Poli A, Olivieri M, et al. Seroprevalence of *Chlamydia pneumoniae* antibodies in a young adult population sample living in Verona. *European Community Respiratory Health Survey (ECRHS) Verona. Infection* 2000; 28: 38-41.
10. Freidank HM, Brauer D. Prevalence of antibodies to *Chlamydia pneumoniae* TWAR in a group of German medical students. *J Infect* 1993; 27: 89-93.
11. Miyashita N, Niki Y, Nakajima M, et al. Prevalence of asymptomatic infection with *Chlamydia pneumoniae* in subjectively healthy adults. *Chest* 2001; 119: 1416-9.
12. Grayston JT, Aldous MB, Easton A, et al. Evidence that *Chlamydia pneumoniae* causes pneumonia and bronchitis. *J Infect Dis* 2001; 168: 1231-5.
13. Kuo CC, Jackson LA, Campbell LA, et al. *Chlamydia pneumoniae* (TWAR). *Clin Microbiol Rev* 1995; 8: 451-61.
14. Bauriedel G, Andrie R, Likungu JA, et al. Persistence of *Chlamydia pneumoniae* in coronary plaque tissue. A contribution to infection and immune hypothesis in unstable angina pectoris. *Dtsch Med Wochenschr* 1999; 124: 1408-13.
15. Fortini A, Corti G, Cappelletti C. *Chlamydia pneumoniae* and atherosclerosis. *Ann Ital Med Int* 1999; 14: 253-63.
16. Balin BJHC, Gerard EJ, Arking DM, et al. Whittum-Hudson Identification and localization of *Chlamydia pneumoniae* in the Alzheimer's brain. *Med Microbiol Immunol* 1998; 187: 23-42.
17. Budak F, Vahaboglu H. The investigation of *Chlamydomphila pneumoniae* in patients with multiple sclerosis. *Int J Neurosci* 2007; 117: 409-15.
18. West SK, Kohlhepp SJ, Jin R, et al. Detection of circulating *Chlamydomphila pneumoniae* in patients with coronary artery disease and healthy control subjects. *Clin Infect Dis* 2009; 48: 560-7.
19. Rodel J, Woytas M, Groh A, et al. Production of basic fibroblast growth factor and interleukin 6 by human smooth muscle cells following infection with *Chlamydia pneumoniae*. *Infect Immun* 2000; 68: 3635-41.
20. Kol A, Bourcier T, Lichtman AH, et al. Chlamydial and human heat shock protein 60s activate human vascular endothelium, smooth muscle cells and macrophages. *J Clin Invest* 1999; 103: 571-7.
21. Kaukoranta-Tolvanen SS, Ronni T, Leinonen M, et al. Expression of adhesion molecules on endothelial cells stimulated by *Chlamydia pneumoniae*. *Microb Pathog* 1996; 21: 407-11.
22. Caspar-Bauguil S, Puissant B, Nazzari D, et al. *Chlamydia pneumoniae* induces interleukin-10 production that down-regulates major histocompatibility complex class I expression. *J Infect Dis* 2000; 182: 1394-401.

23. Geng Y, Shane RB, Berencsi K, et al. Chlamydia pneumoniae inhibits apoptosis in human peripheral blood mononuclear cells through induction of IL-10. *J Immunol* 2000; 164: 5522-9.
24. Byrne GI, Ojcius DM. Chlamydia and apoptosis: life and death decisions of an intracellular pathogen. *Nat Rev Microbiol* 2004; 2: 802-8.
25. Sarris AH, Esgleyes-Ribot T, Crow M, et al. Cytokine loops involving interferon-gamma and IP-10, a cytokine chemotactic for CD4+ lymphocytes: an explanation for the epidermotropism of cutaneous T-cell lymphoma? *Blood* 1995; 86: 651-8.
26. Abrams JT, Vonderheid EC, Kolbe S, et al. Sezary T-cell activating factor is Chlamydia pneumoniae-associated protein. *Clin Diagn Lab Immunol* 1999; 6: 895-905.
27. Halme S, Saikku P, Surcel HM. Characterization of Chlamydia pneumoniae antigens using human T cell lines. *Scand J Immunol* 1997; 45: 378-84.
28. Yawalkar N, Ferenczi K, Jones DA. Profound loss of T-cell receptor repertoire complexity in cutaneous T-cell lymphoma. *Blood* 2003; 102: 4059-66.
29. Hahtola S, Tuomela S, Elo L, et al. Th1 response and cytotoxicity genes are down-regulated in cutaneous T-cell lymphoma. *Clin Cancer Res* 2006; 12: 4812-21.
30. Okamura H, Tsutsi H, Komatsu T, et al. Cloning of a new cytokine that induces IFN-gamma production by T cells. *Nature* 1995; 378: 88-91.
31. Naik SM, Cannon G, Burbach GJ, et al. Human keratinocytes constitutively express interleukin-18 and secrete biologically active interleukin-18 after treatment with pro-inflammatory mediators and dinitrochlorobenzene. *J Invest Dermatol* 1999; 113: 766-72.
32. Yoshimoto T, Takeda K, Tanaka T, et al. IL-12 up-regulates IL-18 receptor expression on T cells, Th1 cells, and B cells: synergism with IL-18 for IFN-gamma production. *J Immunol* 1998; 161: 3400-7.
33. El-Mezayen RE, Matsumoto T. In vitro responsiveness to IL-18 in combination with IL-12 or IL-2 by PBMC from patients with bronchial asthma and atopic dermatitis. *Clin Immunol* 2004; 111: 61-8.
34. Rook AH, Wood GS, Yoo EK, et al. Interleukin-12 therapy of cutaneous T-cell lymphoma induces lesion regression and cytotoxic T-cell responses. *Blood* 1999; 94: 902-8.
35. Yoshimoto T, Tsutsui H, Tominaga K, et al. IL-18, although antiallergic when administered with IL-12, stimulates IL-4 and histamine release by basophils. *Proc Natl Acad Sci USA* 1999; 96: 13962-6.
36. Hwang ST, Janik JE, Jaffe ES, et al. Mycosis fungoides and Sezary syndrome. *Lancet* 2008; 371: 945-57.
37. Adams AE, Zwicker J, Curiel C. Aggressive cutaneous T-cell lymphomas after TNFalpha blockade. *J Am Acad Dermatol* 2004; 51: 660-2.
38. Kaplan EH, Rosen ST, Norris DB, et al. Phase II study of recombinant human interferon gamma for treatment of cutaneous T-cell lymphoma. *J Natl Cancer Inst* 1990; 82: 208-12.
39. Laroche L, Kaiserlian D. Decreased natural-killer-cell activity in cutaneous T-cell lymphomas. *N Engl J Med* 1983; 308: 101-2.
40. Vowels BR, Cassin M, Vonderheid EC, et al. Aberrant cytokine production by Sezary syndrome patients: cytokine secretion pattern resembles murine Th2 cells. *J Invest Dermatol* 1992; 99: 90-4.
41. Wysocka M, Zaki MH, French LE, et al. Sezary syndrome patients demonstrate a defect in dendritic cell populations: effects of CD40 ligand and treatment with GM-CSF on dendritic cell numbers and the production of cytokines. *Blood* 2002; 100: 3287-94.
42. Axelrod PI, Lorber B, Vonderheid EC. Infections complicating mycosis fungoides and Sezary syndrome. *JAMA* 1992; 267: 1354-8.
43. Lee J, Richardson S, Melhem ER, et al. Progressive multifocal leukoencephalopathy from JC virus in a patient with advanced mycosis fungoides. *J Am Acad Dermatol* 2007; 57: 893-5.
44. Park CS, Kim TB, Moon KA, et al. Chlamydomphila pneumoniae enhances secretion of VEGF, TGF-b, and TIMP-1 from human bronchial epithelial cells under Th2 dominant microenvironment. *Allergy Asthma Immunol Res* 2010; 2: 41-7.
45. Rössler MJ, Rappl G, Muche M, et al. No evidence of skin infection with Chlamydia pneumoniae in patients with cutaneous T cell lymphoma. *Clin Microbiol Infection* 2003; 9: 721-3.
46. Ferreri AJ, Ponzoni M, Govi S, et al. Prevalence of chlamydial infection in a series of 108 primary cutaneous lymphomas. *Br J Dermatol* 2012; 166: 1121-3.
47. Caselli E, Borghi A, Maritati M, et al. Relapses of primary cutaneous anaplastic large-cell lymphoma in a female immunocompetent patient with persistent Chlamydomphila pneumoniae and human herpesvirus 8 infection. *Infect Agent Cancer* 2016; 11: 31.